

<https://helda.helsinki.fi>

Methanotrophy induces nitrogen fixation during peatland development.

Larmola, Tuula

2014

Larmola , T , Leppänen , S , Tuittila , E-S , Aarva , M , Merilä , P , Fritze , H & Tirola , M
2014 , ' Methanotrophy induces nitrogen fixation during peatland development. ' ,
Proceedings of the National Academy of Sciences of the United States of America , vol. 111
, no. 2 , pp. 734-739 . <https://doi.org/10.1073/pnas.1314284111>

<http://hdl.handle.net/10138/224708>

<https://doi.org/10.1073/pnas.1314284111>

cc_by_nc_nd

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Methanotrophy induces nitrogen fixation during peatland development

Tuula Larmola^{a,1}, Sanna M. Leppänen^b, Eeva-Stiina Tuittila^{a,2}, Maija Aarva^b, Päivi Merilä^c, Hannu Fritze^d, and Marja Tiirola^b

^aForest Sciences, University of Helsinki, FI-00014, Helsinki, Finland; ^bBiological and Environmental Science, University of Jyväskylä, FI-40014, Jyväskylä, Finland; ^cFinnish Forest Research Institute, University of Oulu, FI-90014, Oulu, Finland; and ^dFinnish Forest Research Institute, FI-01301, Vantaa, Finland

Edited by Sarah Hobbie, University of Minnesota, Saint Paul, MN, and approved November 26, 2013 (received for review July 28, 2013)

Nitrogen (N) accumulation rates in peatland ecosystems indicate significant biological atmospheric N₂ fixation associated with *Sphagnum* mosses. Here, we show that the linkage between methanotrophic carbon cycling and N₂ fixation may constitute an important mechanism in the rapid accumulation of N during the primary succession of peatlands. In our experimental stable isotope enrichment study, previously overlooked methane-induced N₂ fixation explained more than one-third of the new N input in the younger peatland stages, where the highest N₂ fixation rates and highest methane oxidation activities co-occurred in the water-submerged moss vegetation.

CH₄ | diazotrophy | mire | peat | phosphorus

Peat-accumulating wetlands, i.e., peatlands, store approximately 30% of the global soil carbon (C) (1), and this value is even higher if the permafrost regions in the Northern Hemisphere are also taken into account (2). As peat accumulates, the ecosystem becomes independent of the groundwater influence and the vegetation becomes more nutrient limited. This gradual succession from minerotrophic fen to *Sphagnum*-dominated ombrotrophic bog ecosystem is the general peatland development pattern. Because the growth and decomposition rates of *Sphagnum* mosses are greatly responsible for the C biosequestration in peatlands, the ecology of *Sphagnum* mosses is of particular interest. Recent studies have shown that *Sphagnum* mosses have an association with methanotrophic bacteria that leads to a reduction in methane (CH₄) emissions to the atmosphere and the provision of additional carbon dioxide (CO₂) source for the host plants (3, 4). However, the growth of *Sphagnum* mosses in peatlands is often N limited, at least under low atmospheric N deposition (5), so biological fixation of atmospheric N₂, i.e., the biological conversion of dinitrogen to plant-available ammonium, may stimulate moss growth (6). Under low atmospheric N deposition, moss-associated cyanobacteria have been shown to play an important role in the N budget in forests (7, 8) and peatlands (9), where N₂ fixation is favored by moist conditions. In peatlands, however, molecular analyses of genes that encode nitrogenase reductase proteins (*nifH*) in *Sphagnum* mosses have indicated that moss-associated N₂ fixers (diazotrophs) belong mainly to the metabolically diverse class Alphaproteobacteria (10), which includes phototrophic, heterotrophic, and methanotrophic genera.

In previous studies, N₂ fixation rates in peatlands have been found to correlate with the minerotrophy of peatlands, in particular with the level of phosphorus (P) (11), but the nutrient controls of N₂ fixation have not been linked to peatland succession toward the bog stage. Further, in these studies, N₂ fixation has been measured by using acetylene reduction assay (12), which does not provide a quantitative measure of N added to the system, because it inhibits the activity of many noncyanobacterial diazotrophs, but specifically methanotrophic bacteria by inactivating the essential methane monooxygenase enzyme (13). This inhibition is a serious drawback, because a broad range of methanotrophic bacteria contains genes that code for the N₂ fixation pathway and shows nitrogenase activity (11, 14, 15). Because of this methodological problem, the role

of methanotrophic N₂ fixation and the relationship of N₂ fixation with C cycling have not previously been evaluated at an ecosystem scale. The elucidation of the linkage between methanotrophy and the overall N cycle in peatlands becomes feasible by the application of stable isotope (¹⁵N₂) techniques (8, 16).

We studied N₂ fixation and CH₄ oxidation in the dominant flark and hummock vegetation of 12 pristine peatlands, which varied in age from 200 to 2,500 y due to still ongoing postglacial rebound on the coast of Bothnian Bay, Finland (Fig. 1 and Table S1). Together with the Hudson Bay Lowlands in Canada, the Bothnian Bay of the Baltic Sea between Finland and Sweden is the region where the rebound after the pressure of ice mass and the consequent formation of new land from the sea is most rapid. This chronosequence of peatlands offers an exceptional opportunity to study the links between N and C cycling over an undisturbed peatland gradient.

In young peatlands, unstable hydrological conditions are likely to result in low CH₄ emissions and, thereby, small methanotrophic communities (17). As the hydrological conditions become more stable in the later successional stages with increasing peat depths, CH₄ emissions increase (18). High CH₄ emissions have been linked to minerotrophic fen stages, in which the dense sedge vegetation readily provides substrate for methanogenesis, whereas in ombrotrophic bogs characterized by *Sphagnum* mosses and dwarf shrubs, the rates of decomposition and substrate supply for the CH₄ production are slow (19). However, the ranges for *Sphagnum* hosted

Significance

In peatlands, the external sources of nitrogen are mainly atmospheric, but the atmospheric nitrogen deposition alone cannot explain the long-term annual nitrogen accumulation rates to these ecosystems. Because of methodological problems, methane-induced fixation of atmospheric dinitrogen gas has been previously overlooked as an additional nitrogen input mechanism. We found that the activity of methane-oxidizing bacteria provides not only carbon but also nitrogen to peat mosses and, thus, contributes to carbon and nitrogen accumulation in peatlands, which store approximately one-third of the global soil carbon pool. Our results imply that nitrogen fixation in wetlands may be strongly underestimated when methods inhibiting methane oxidizers are used.

Author contributions: T.L., S.M.L., E.-S.T., H.F., and M.T. designed research; T.L., S.M.L., and M.A. performed research; S.M.L., E.-S.T., P.M., and M.T. contributed new reagents/analytic tools; T.L. analyzed data; and T.L., E.-S.T., and M.T. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed at the present address. Environmental Studies Department, Mount Holyoke College, South Hadley, MA 01075. E-mail: tuula.larmola@gmail.com.

²Present address: School of Forest Sciences, University of Eastern Finland, FI-80101, Joensuu, Finland.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1314284111/-DCSupplemental.




Fig. 1. Carbon (left axis downward) and nitrogen (right axis downward) accumulation rates during the 2,500-y primary succession of Siikajoki peatlands from meadows, mesotrophic fens, and oligotrophic fens toward fen-bog transitions on the land-uplift coast of the Bothnian Bay (age cohorts SJ2–SJ5; ref. 23). The error bars are SD. The estimates of site age means are based on the land-uplift rate (meadows and mesotrophic fens) or on radiocarbon dating (oligotrophic fens and fen-bog transitions); m a.s.l., meters above sea level; BP, before present.

CH_4 oxidation rates in minerotrophic fens and ombrotrophic bogs have been found to overlap (4). It has been hypothesized that despite the low inorganic N concentration in bogs, CH_4 oxidation is not N limited because methanotrophic N_2 fixation may compensate for the N requirement of methanotrophic bacteria (20).

With reference to these factors, we hypothesized that (i) CH_4 induces moss-associated N_2 fixation, which would be most pronounced in the late stages of peatland succession and (ii) the overall N_2 fixation rates along the peatland succession gradient are governed by gradually changing environmental factors, such as water table depth, availability of CH_4 in the pore water, and concentrations of P, iron (Fe), or molybdenum (Mo). The rationale for studying the role of Fe and Mo was that the two methane monooxygenases (e.g., ref. 21) and all three nitrogenase isoenzymes contain Fe, and one of the nitrogenase isoenzymes contains Mo as a necessary cofactor (22). We tested the two hypotheses in a $^{15}\text{N}_2$ and $^{13}\text{CH}_4$ pulse-labeling experiment, where moss samples collected from flark and hummock habitats of four successional peatland stages were incubated in situ with and without CH_4 addition, each under prevailing light conditions and in the dark. These treatments were used to reveal whether N_2 fixation was attributed to photosynthetic, heterotrophic, or methanotrophic activity.

Results and Discussion

Fixation of N_2 was found in the live parts of the *Sphagnum* mosses in all of the studied moss patches. Although *Sphagnum*-associated N_2 fixation was observed in all of the study peatlands of different successional stages, the process was significantly higher (five- to ninefold) in the wet depressions (flarks) of the midsuccessional mesotrophic fens compared with the other successional stages, which consisted of younger meadows, older oligotrophic fens and fen-bog transitions (Fig. 2A and Tables S2 and S3). When the summertime N_2 fixation rates ($0\text{--}126\text{ nmol}\cdot\text{g}^{-1}$ of moss biomass $\cdot\text{h}^{-1}$) were converted to annual areal values ($0.1\text{--}2.9\text{ g of N}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$; Fig. 2B), they were up to 10 times greater than the current inorganic N deposition rates ($0.3\text{ g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$) for the region. These annual values indicate that *Sphagnum*-associated N_2 fixation is a major N input to boreal peatlands and, in conjunction with atmospheric N deposition, explains the long-term annual N accumulation rate ($0.6\text{--}1.3\text{ g}\cdot\text{m}^{-2}$; ref. 23) in the studied peatland ecosystems (Fig. 2B). Our areal estimations for fen stages and fen-bog transition are within the wide range of the admittedly few estimates available for *Sphagnum*-associated N_2 fixation in temperate, boreal, and subarctic ecosystems. These range from $0.1\text{--}6.4\text{ g of N}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ for fens (11, 16, 24) and $0.1\text{--}1\text{ g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$ for bogs (9, 25). Nevertheless, our estimates may be rough because of the large SEs and the extrapolation over time: We assume a 6-mo active season, but substantial heterotrophic

microbial activity in addition to phototrophic activity could occur during winter. At these sites, 30–65% of annual moss growth occurs during the October–April period, outside the traditionally defined growing season (26).

In all studied peatland stages except wet meadows, incubation under prevailing light conditions resulted in enhanced N_2 fixation, on average threefold in comparison with the dark treatments, which suggests that phototrophic organisms may be the most active N_2 fixers or that photosynthesis provides carbohydrates to fuel heterotrophic N_2 fixers (Fig. 2A). Methane-induced N_2 fixation contributed approximately 40% (33–47%) of the N_2 fixation in the three younger peatland stages, but was negligible in the fen-bog transition stage (Fig. 2A and B). Thus, the hypothesis of a larger methanotrophic contribution to N_2 fixation in late successional stages was not supported. The rate of the biomass incorporation of $^{13}\text{CH}_4$ -derived C indicated that moss-associated methanotrophy was also highest in mesotrophic fens and continued at moderate rates in the flarks of the older stages (Fig. 2C), where CH_4 addition did not enhance N_2 fixation. The meadows showed significantly lower CH_4 oxidation rates compared with the other stages (Table S2). A comparison of the successional patterns of N accumulation and N_2 fixation (Fig. 2B) provides further evidence of the ecological and biogeochemical importance of N_2 fixation and methanotrophic N_2 fixation during peatland development. Based on peat profiles from our sites, 70% of N and 40% of C accumulated during the first 1,000 y of the 2,500-y period (23), a period during which N_2 fixation peaked and had the strongest response to CH_4 addition. This pattern implies that methanotrophic N_2 fixation contributes to rapid N accumulation in the fen stages. The predicted warming conditions in the northern latitudes (27) may impact on boreal peatland development in two ways: At the southern limit of permafrost, melting is promoting a reverse succession from ombrotrophic bog to fen ecosystem, whereas at the southern border of the fen region (aapamires, wet fen-dominated peatland complexes) drainage due to increased evapotranspiration may accelerate ombrotrophication, i.e., succession toward bog ecosystem (28). Our results indicate that these successional changes are likely to lead to changes in N_2 dynamics.

Based on the growth rates of *Sphagnum* species at our sites (26) and site-specific N content of the *Sphagnum* species (Table S4), we estimate that N obtained by N_2 fixation could correspond to an average $37 \pm 18\%$ (mean \pm SEM, range 4–58%, $n = 6$) of the moss biomass N increment. This proportion is in agreement with the recent estimate of 35% for *Sphagnum riparium*, inferred from N content and growth rate in a 2-mo laboratory experiment (6). Despite the low absolute areal rate of N_2 fixation in the older mainly rainwater-fed stages (fen-bog transitions), where moss growth is more nutrient limited, the comparison of *Sphagnum*




Fig. 2. *Sphagnum* moss-associated N_2 fixation and CH_4 oxidation (biomass incorporation of CH_4 -derived C) in 12 peatlands of the peatland chronosequence (meadows, mesotrophic and oligotrophic fens, and fen-bog transitions) based on the stable isotope pulse-labeling experiment. The error bars are SEM ($n = 3$ peatlands in each stage). (A) Contribution of dark (heterotrophic, black), light-induced (phototrophic, white), and CH_4 -induced (methanotrophic, cross-hatched) N_2 fixation in flark (F) and hummock (H) vegetation. The meadows have only flark *Sphagnum*. (B) The contribution of CH_4 -induced N_2 fixation (cross-hatched) to *Sphagnum*-associated N_2 fixation.

growth rates (26) and N contents further suggests that the proportion of fixed N of the new biomass N increment may increase to 58%. The rest of the N is, we presume, being taken up as inorganic ions and organic N, or recycled to new growth from older parts of the moss shoot (29). The time scale at which N fixed in the moss becomes available for the moss host and for other plants may range from fast exchange over a time span of days (6, 11) to slower nutrient release from decomposing *Sphagnum* litter over a period of years.

Our results showed faster CH_4 -C biomass incorporation in light than in dark (Fig. 2C), which suggests that mosses fixed additional CH_4 -derived CO_2 during photosynthesis. The contribution of CH_4 -derived C was 26% in light, but 10% in dark (calculated as above for N) for *Sphagnum* C in the flarks. In hummocks, where CH_4 and CO_2 concentrations are at atmospheric levels, CH_4 -derived C contributed to only 0–3% of the incorporated C in light and dark. These findings indicate that CH_4 can be a significant C source for submerged *Sphagnum*, supporting the results of previous studies (3, 4).

The moss ratios of N:P, C:N, and C:P, integrated indices of nutrient availability, showed decreasing ratios with increasing N_2 fixation activity (Spearman ρ –0.32, –0.61, and –0.54, for N:P, C:N, and C:P, respectively, $P < 0.003$). The successional pattern in N_2 fixation rates was best explained by P availability (Fig. 3 and Tables S4 and S5). The highest rates of N_2 fixation were associated with the lowest moss N:P ratios, being lower than the threshold for N limitation in soil microbes (N:P < 7:1; ref. 30). Thus, N_2 fixers were able to respond to N demand relative to P supply, as shown (e.g., ref. 31). In the older stages, N:P ratios indicated that plant growth was limited by P (32), yet active moss associated N_2 fixation persisted there as well (Figs. 2 and 3).

Although the moss P content was the sole significant predictor of N_2 fixation in the light treatment without CH_4 , the contribution of CH_4 -induced N_2 fixation was best explained by the combination of the moss P and Fe contents (Table S5). Among the sampled microhabitats, moss Fe content correlated strongly with the water table depth below the moss capitula and pH (Table S4). Thus, provided that P supply was sufficient, CH_4 -induced N_2 fixation correlated with CH_4 oxidation (hypothesis 2; Fig. 3), but the rate of CH_4 oxidation as such was not the primary factor controlling CH_4 -induced N_2 fixation in these sites. The CH_4 oxidation was, in turn, best explained by the moss Fe content that depended on the water table level (Fig. S1). Concurrent CH_4 concentration in pore water did not significantly correlate with N_2 fixation (Spearman rank correlation $P = 0.96$, $n = 84$) or biomass incorporation of CH_4 -derived C ($P = 0.16$, $n = 42$).

Overall, our results demonstrate that methanotrophy and N_2 fixation are tightly linked in the wet fen depressions and that methanotrophic activity enhances N_2 fixation (hypothesis 1). However, because the interactions within the endosymbiotic microbial communities complicate the analysis, we do not directly interpret the experimentally observed dark, light-induced, and CH_4 -induced rates as heterotrophic, phototrophic, and methanotrophic N_2 fixation. The experimental approach cannot distinguish whether the CH_4 -induced N_2 fixation equals methanotrophic N_2

Data are averages of light and dark incubations in $^{15}N_2$ and $^{15}N_2 + ^{13}CH_4$ treatments, respectively, weighted with the proportions of flark and hummock microhabitats in each successional stage. Long-term average peat N accumulation rates for the sites (total accumulation divided by the year since peatland initiation; ref. 23) is shown with filled circles. (C) Incorporation of $^{13}CH_4$ -C into the biomass (moss + microbes) based on the weighted averages of flark and hummock microhabitats in each successional stage. In each bar, the filled area indicates the incorporation of $^{13}CH_4$ -derived C in the dark (i.e., incorporation of CH_4 into methanotroph biomass) and the open area indicates the average additional incorporation of $^{13}CH_4$ -derived C under prevailing light conditions (i.e., incorporation of CO_2 emitted by methanotrophs into autotrophic plant or microbial biomass via photosynthesis).




Fig. 3. The relationships between the rates of N_2 fixation, CH_4 oxidation and phosphorus content in *Sphagnum* moss samples of different successional stages of peatlands (●, meadow; ■, mesotrophic fen; ▲, oligotrophic fen, and ▼, fen-bog transition). (A) The relationship between N_2 fixation (\ln transformed) and moss P content ($\ln N_2$ fixation = $1.25 \pm 0.14[\text{P content}] + 0.36 \pm 0.13$ (coefficient \pm SE), $R^2 = 0.49$, $\text{df}_{\text{reg, res}} = 1, 82$; Table S5). The moss N:P ratios > 16 indicating P limitation (32) are shown with open symbols. Averages for all four treatments per microhabitat and stage are shown ($n = 84$ incubations). (B) The Spearman rank correlation between the rates of CH_4 oxidation (measured as biomass incorporation of CH_4 -derived C) and CH_4 -induced N_2 fixation (calculated as the difference in N_2 fixation between $^{15}\text{N}_2 + ^{13}\text{CH}_4$ and $^{15}\text{N}_2$ only treatments). The moss P content $< 0.5 \text{ mg g}^{-1}$ is indicated by open symbols ($n = 42$ incubations).

fixation, because the observed CH_4 induction may be, at least partly, indirect. The higher CO_2 concentrations that result from CH_4 oxidation inside the *Sphagnum* cells would enhance photosynthetic N_2 fixation (Fig. 4). Cell-level identification of the correct nutritional mode of the N_2 fixing organisms may require single cell analysis, which is possible by using secondary ion mass spectrometry. Similarly, dark and light-induced N_2 fixation may not equal phototrophic and heterotrophic N_2 fixation, because light can increase the cellular concentration of C substrates and oxygen, which are to some extent essential to N_2 fixation of aerobic heterotrophs.

In all, our findings imply that interlinks of N_2 fixation and CH_4 cycling constitute an important mechanism that fulfils the need of new N and explains the rapid peat and N accumulation rates during the fen stages of peatland development. Our results

further support the idea that N_2 fixation is the primary mechanism linking N and P availability, as suggested for terrestrial and aquatic ecosystems (e.g., refs. 31 and 33). Both the N_2 fixation rates and the methanotrophic contribution to N_2 fixation along the peatland succession gradient were primarily governed by P availability. The successional patterns of different microbial groups involved in N_2 fixation were further regulated by water table level and Fe content, which controlled both the N_2 fixation and CH_4 oxidation activities of the plant-associated bacteria. In conclusion, moss-associated microbes may explain the dominance of *Sphagnum* mosses in nutrient-poor peatlands: besides providing an additional in-cell CO_2 source for photosynthesis, these microbial communities provide new N in the cold and acidic environment where decomposition and nutrient recycling are otherwise slow.

Materials and Methods

Field Site. The Siikajoki peatland chronosequence is a replicated primary successional series of peatland ecosystems in the land-uplift coast of Bothnian Bay, Finland ($64^\circ 45' \text{N}$, $24^\circ 42' \text{E}$; refs. 23 and 34). Our study sites at Siikajoki consisted of 12 peatlands that represent four successional stages, which vary in age from 200 to 2,500 y (Fig. 1). Each stage had three replicates (Table S1). The sites were located within 8 km on the coast of Bothnian Bay, where the rate of land uplift is approximately 8 mm y^{-1} due to ongoing postglacial rebound (35). The mean annual temperature is 2.3°C , and the mean annual precipitation 521 mm. The inorganic N deposition rate in the region is $0.3 \text{ g m}^{-2} \text{ y}^{-1}$ (36).

The peatlands, 0.5–1.5 ha in size, have developed in depressions between sand dune formations with similar underlying soil (18, 37, 38). Along the chronosequence, the vegetation changes from flooded *Carex nigra* and *Agrostis canina* dominated meadows in the early stages to *Carex*-dominated mesotrophic and oligotrophic fens and to fen-bog transitions where the bog vegetation is characterized by an increasing cover of dwarf shrubs in hummock surfaces. The moss layer in the meadows is sparse (total moss cover $< 20\%$) and dominated by *Warnstorfia* spp. mosses, with a *Sphagnum* cover of approximately 2%. In the fen and bog stages, *Sphagnum* mosses cover 50–70% of the peat surface (total moss cover 55–90%). The fen stages show a distinct development of microtopography with *Sphagnum subsecundum* and *Sphagnum majus* dominated flarks (wet depressions) followed by *Sphagnum fimbriatum* and then *Sphagnum papillosum* dominated hummocks which lead in the oldest stages to colonization by *Sphagnum fuscum*, a species characteristic of bog hummocks.




Fig. 4. The interplay of N_2 -fixing bacteria in the upper moss vegetation. Phototrophic, heterotrophic, and methanotrophic bacteria all take part in the fixation of atmospheric N_2 to plant available N (NH_4^+). The supply of CH_4 may support the direct methanotrophic N_2 fixation, but it may also provide additional CO_2 for phototrophic microbes, as well as for the moss itself. Both phototrophs and methanotrophs provide organic molecules (CH_2O) to heterotrophic N_2 fixers. Methanotrophic activity may also decrease the oxygen tension, providing a more suitable microenvironment for the oxygen-sensitive nitrogenase enzyme complex.

Field Experiment. In early June 2010, we collected the dominant *Sphagnum* mosses from both flark and hummock habitats at each site. Moss sampling was based on a vegetation survey made earlier at the sites (23). We incubated the moss samples in situ for 2 d under three treatments (A $^{15}\text{N}_2 + ^{13}\text{CH}_4$, B $^{15}\text{N}_2$, and C unlabeled control), each under prevailing light conditions (day length 20 h) and in the dark. We hypothesized that in the $^{15}\text{N}_2 + ^{13}\text{CH}_4$ treatment under prevailing light conditions, all potential diazotrophs (cyanobacteria, heterotrophs, methanotrophs) can be active. In the $^{15}\text{N}_2$ treatment under prevailing light, cyanobacteria and heterotrophs can actively fix N_2 . The $^{15}\text{N}_2$ treatment in the dark will provide information on the activity of heterotrophs other than obligate methanotrophs. Additionally, the $^{13}\text{CH}_4$ treatments will determine the incorporation of CH_4 -derived C in to the microbial biomass. Under prevailing light, this treatment will also show the potential of mosses to use CH_4 -derived CO_2 in photosynthesis. Mosses (1 g in dry mass) were incubated in situ in 120-mL glass vials. Only the upper, live parts were used in the incubation, and 10 mL of water from the sampling site was added into the vials to keep the samples moist. A volume of 20 mL of air was removed, and 20 mL of $^{15}\text{N}_2$ tracer gas [98% (vol/vol) enriched; Cambridge Isotope Laboratories] and 1.2 mL of $^{13}\text{CH}_4$ tracer gas [99% (vol/vol) enriched; CK Gas Products] were injected into the vials to reach headspace enrichment levels of 21% (vol/vol) and 99% (vol/vol) for $^{15}\text{N}_2$ and $^{13}\text{CH}_4$, respectively. Control samples were incubated in the ambient air, and the dark-treated samples were covered with aluminum foil. The vials were placed where the samples were collected, so that half of the vial remained above the water level. Incubations were terminated after 45 h by opening the vials, emptying them of water, and freezing them at -20°C . Then the samples were dried at $+50^\circ\text{C}$ to a constant mass and analyzed for their bulk $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values and C and N contents using a stable isotope ratio mass spectrometer as described in ref. 8. The internal precision (SD of replicate standards) of the isotope analysis was always better than 1.5‰ for $\delta^{15}\text{N}$ and 0.1 for $\delta^{13}\text{C}$.

During the field incubations, the temperature in the moss layer (measured by using iButton; Maxim) averaged 14.7°C , with a range of $5\text{--}30^\circ\text{C}$, and did not differ significantly among the sites. In each microhabitat, we measured the water table depth within a pipe well and drew pore water samples to determine pH and dissolved CH_4 concentrations. The CH_4 concentration in water samples of 10 mL was determined by using the headspace equilibration technique (39) by following ref. 4. The concentration of CH_4 was analyzed in the gas headspace by using a gas chromatograph equipped with flame ionization detector (Agilent 7890A; Agilent Technologies). Dissolved CH_4 concentrations were calculated from headspace concentrations according to Henry's law by using the values after ref. 40. We took

volumetric samples of each moss species from the uppermost 10 cm to determine density. After removing debris and other mosses, volumetric samples of *Sphagnum* were dried at 60°C for 48 h and weighed to convert the incubation results to N_2 fixation and CH_4 -C biomass incorporation (CH_4 oxidation) rates per peatland surface area (m^2). A subset of plots ($n = 1$ in each stage) was surveyed for the relative proportions of flark and hummock habitats by measuring the water table depth in a network of 8–12 wells. Phosphorus, Fe, and Mo (below detection limit) contents in the tissue of the incubated *Sphagnum* samples were extracted with HNO_3 by using wet digestion (EPA-3051) and measured by using a plasma emission spectrometer (ICP-OES; IRIS Intrepid II XSP).

Data Analyses. The rates of N_2 fixation and CH_4 oxidation measured as CH_4 -C incorporation in to the moss biomass were calculated as enrichment values and thereafter as fixation rates ($\text{nmol}\cdot\text{g}^{-1}$ of dry moss $\cdot\text{h}^{-1}$) by following ref. 8. The values represent the actual ^{15}N and ^{13}C uptake in each sample during the 45-h incubation. The rates of N_2 fixation, CH_4 -induced N_2 fixation, and CH_4 -C biomass incorporation were natural-log (ln) transformed to meet the requirements of normality. The effects of terrestrial age and treatments on N_2 fixation rates were analyzed with four-way nested analysis of covariance (ANCOVA) with successional stage, site nested within the successional stage, light treatment, and CH_4 treatment as the four factors. Water table depth was used as a covariate to take in to account microtopography. The effects of terrestrial age and light on the rates of biomass incorporation of CH_4 -derived C were analyzed with three-way nested ANCOVA with successional stage, site nested within the successional stage and light treatment as the three factors and water table depth as a covariate. Pair-wise differences in successional stages were tested by using Bonferroni post hoc tests. The relationships between environmental variables (nutrient contents, water table depth, water pH, and CH_4 concentration) and N_2 fixation, CH_4 -induced N_2 fixation, and the biomass incorporation of CH_4 -derived C were analyzed by using Spearman rank correlations and stepwise regressions.

ACKNOWLEDGMENTS. We thank L. Kuusisto for nutrient analyses, T. Sinisalo for help in the stable isotope mass spectrometer analyses, I. Murtovaara for designing the schematic figures, and D. Wilson for English revision. Funding was provided by Academy of Finland Projects 121535 (to T.L.), 131409 and 218101 (to E.-S.T.), and 260797 (to M.T.), and by Biological Interactions Graduate School (S.M.L.).

- Gorham E (1991) Northern peatlands: Role in the carbon cycle and probable responses to climatic warming. *Ecol Appl* 1(2):182–195.
- Tarnocai C, et al. (2009) Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochem Cycles* 23: 10.1029/2008GB003327.
- Raghoebarsing AA, et al. (2005) Methanotrophic symbionts provide carbon for photosynthesis in peat bogs. *Nature* 436(7054):1153–1156.
- Larmola T, et al. (2010) The role of *Sphagnum* mosses in the methane cycling of a boreal mire. *Ecology* 91(8):2356–2365.
- Aerts R, Wallen B, Malmer N (1992) Growth-limiting nutrients in *Sphagnum*-dominated bogs subject to low and high atmospheric nitrogen supply. *J Ecol* 80(1): 131–140.
- Berg A, Danielsson Å, Svensson BH (2013) Transfer of fixed-N from N_2 -fixing cyanobacteria associated with the moss *Sphagnum* riparium results in enhanced growth of the moss. *Plant Soil* 362(1–2):271–278.
- DeLuca TH, Zackrisson O, Nilsson M-C, Sellstedt A (2002) Quantifying nitrogen-fixation in feather moss carpets of boreal forests. *Nature* 419(6910):917–920.
- Leppänen SM, Salemaa M, Smolander A, Mäkipää R, Tirola M (2013) Nitrogen fixation in forest mosses along a N deposition gradient. *Environ Exp Bot* 90:62–69.
- Granhall U, Selander H (1973) Nitrogen fixation in a subarctic mire. *Oikos* 24(1):8–15.
- Bragina A, Berg C, Müller H, Moser D, Berg G (2013) Insights into functional bacterial diversity and its effects on Alpine bog ecosystem functioning. *Sci Rep* 3:1955.
- Basilier K, Granhall U, Stenstrom TA (1978) Nitrogen fixation in wet minerotrophic moss communities of a subarctic mire. *Oikos* 31(2):236–246.
- Hardy RWF, Holsten RD, Jackson EK, Burns RC (1968) The acetylene-ethylene assay for $\text{n}(\text{g})$ fixation: Laboratory and field evaluation. *Plant Physiol* 43(8):1185–1207.
- Flett RJ, Rudd JWM, Hamilton RD (1975) Acetylene reduction assays for nitrogen fixation in freshwaters: A note of caution. *Appl Microbiol* 29(5):580–583.
- Davis JB, Coty VF, Stanley JP (1964) ATMOSPHERIC NITROGEN FIXATION BY METHANE-OXIDIZING BACTERIA. *J Bacteriol* 88(2):468–472.
- Auman AJ, Speake CC, Lidstrom ME (2001) nifH sequences and nitrogen fixation in type I and type II methanotrophs. *Appl Environ Microbiol* 67(9):4009–4016.
- Gavazov KS, Soudzilovskaia NA, van Logtestijn RSP, Braster M, Cornelissen JHC (2010) Isotopic analysis of cyanobacterial nitrogen fixation associated with subarctic lichen and bryophyte species. *Plant Soil* 333(1–2):507–517.
- Juottonen H, et al. (2012) Methane-cycling microbial communities and methane emission in natural and restored peatlands. *Appl Environ Microbiol* 78(17):6386–6389.
- Leppälä M, Oksanen J, Tuittila E-S (2011) Methane flux dynamics during mire succession. *Oecologia* 165:489–499.
- Nykänen H, Alm J, Silvola J, Tolonen K, Martikainen P (1998) Methane fluxes on boreal peatlands of different fertility and the effect of long-term experimental lowering of the water table on flux rates. *Global Biogeochem Cycles* 12(1):53–69.
- Bodelier PLE, Laanbroek HJ (2004) Nitrogen as a regulatory factor of methane oxidation in soils and sediments. *FEMS Microbiol Ecol* 47(3):265–277.
- Chowdhury TR, Dick RP (2013) Ecology of aerobic methanotrophs in controlling methane fluxes from wetlands. *Appl Soil Ecol* 65:8–22.
- Zehr JP, Jenkins BD, Short SM, Steward GF (2003) Nitrogenase gene diversity and microbial community structure: a cross-system comparison. *Environ Microbiol* 5(7):539–554.
- Tuittila E-S, et al. (2013) Wetland chronosequence as a model of peatland development: Vegetation succession, peat and carbon accumulation. *Holocene* 23(1):25–35.
- Waughman GJ, Bellamy DJ (1980) Nitrogen fixation and the nitrogen balance in peatland ecosystems. *Ecology* 61(5):1185–1198.
- Chapman RR, Hemond HF (1982) Dinitrogen fixation by surface peat and *Sphagnum* in an ombrotrophic bog. *Can J Bot* 60(5):538–543.
- Laine AM, Juurola E, Häjek T, Tuittila E-S (2011) *Sphagnum* growth and ecophysiology during mire succession. *Oecologia* 167(4):1115–1125.
- Intergovernmental Panel on Climate Change (2007): *The Physical Science Basis. Working Group I Contribution to the IPCC Fourth Assessment Report*, eds Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (Cambridge Univ Press, Cambridge, UK).
- Tahvanainen T (2011) Abrupt ombrotrophication of a boreal aapa mire triggered by hydrological disturbance in the catchment. *J Ecol* 99(2):404–415.
- Aldous AR (2002) Nitrogen translocation in *Sphagnum* mosses: effects of atmospheric nitrogen deposition. *New Phytol* 156(2):241–253.
- Cleveland CC, Liptzin D (2007) C:N:P stoichiometry in soil: Is there a “Redfield ratio” for the microbial biomass? *Biogeochemistry* 85(3):235–252.
- Houlton BZ, Wang Y-P, Vitousek PM, Field CB (2008) A unifying framework for di-nitrogen fixation in the terrestrial biosphere. *Nature* 454(7202):327–330.
- Reich PB, Oleksyn J (2004) Global patterns of plant leaf N and P in relation to temperature and latitude. *Proc Natl Acad Sci USA* 101(30):11001–11006.
- Schindler DW (1977) Evolution of phosphorus limitation in lakes. *Science* 195(4275): 260–262.
- Merilä P, et al. (2006) Methanogen communities along a primary succession transect of mire ecosystems. *FEMS Microbiol Ecol* 55(2):221–229.
- Ekman M (1996) A consistent map of the postglacial uplift of Fennoscandia. *Terra Nova* 8(2):158–165.

36. Mustajärvi K, et al. (2008) Fluxes of dissolved organic and inorganic nitrogen in relation to stand characteristics and latitude in Scots pine and Norway spruce stands in Finland. *Bor. Environ. Res* 13(suppl.B):3–21.
37. Leppälä M, Kukko-Oja K, Laine J, Tuittila E-S (2008) Seasonal dynamics of CO₂ exchange during primary succession of boreal mires as controlled by phenology of plants. *Ecoscience* 15(4):460–471.
38. Leppälä M, Laine AM, Seväkivi M-L, Tuittila E-S (2011) Differences in CO₂ dynamics between successional mire plant communities during wet and dry summers. *J Veg Sci* 22(2):357–366.
39. McAuliffe CC (1971) GC determination of solutes by multiple phase equilibration. *Chem Technol* 1:46–51.
40. Lide R, Fredrikse HPR, eds (1995) *CRC Handbook of Chemistry and Physics* (CRC, Boca Raton, FL), 76th Ed.